
KCSE TRIAL 2020

BIOLOGY PAPER 2

SECTION A – (40 MARKS)

Answer ALL Questions in This Section In The Spaces Provided.

1. The diagram below shows the mammalian digestive system. Study it carefully and answer the questions that follow.




	Name the parts labeled A and B	(2marks)
	ii) How is the structure labeled A in the diagram adapted to carry ou	t its function. (2marks)
 b)	i) Name the hormone secreted by the walls of the part labeled C	(1mark)
	ii) Explain the role of the hormone in b) (i) above in digestion.	(3marks)

- 2. In human beings a download pointed frontal hairline ("window peak") is a heritable trait from an expression of recessive gene in a somatic cell. Use 'W' for a dominant gene.
 - a) Determine the F1 generation if a homozygous peak male is married to a homozygous frontal hairlined female parent. (4marks)

	b) State two causes of variations.	(2marks)
	c) Name two sex linked genetic disorders that can affect both hur	man females and males. (2marks)
3.	A student set up an experiment using soaked and dry seeds as show	wn below
	Soaked seeds Cotton wool; Thermometer	Dry seeds
a)	State the objective of this experiment	(1mark)
b)	State the observations made in each of the flasks after 24 hours	(2marks)
c)	Account for the observation made in (b) above	(2marks)
d)	Suggest why vacuum flasks were used in this experiment.	(1mark)

Why should the seeds be washed with antiseptic 10% formalin?	(1mark)
Shown below is a section through the mammalian nephron.	
N N N N N N N N N N N N N N N N N N N	
n) Name the structures labeled A and N.	(2marks)
Name the structures labeled A and N. Name all the structures in a nephron which are normally present in a kidney	
Name all the structures in a nephron which are normally present in a kidney	the cortex region
Name all the structures in a nephron which are normally present in a kidney	the cortex region (1 mark)

5. The diagram below represents a food web in a certain ecosystem.



a)	Name i)	the trophic level occupied by each of the following: Caterpillars	(1 mark)					
	ii)	Small insects	(1mark)					

b) From the food web, construct two food chains which end with lizards as tertiary consumer. (2marks)

i) Which organisms have the least biomass in this ecosystem	
ii) Explain the answer in (i) above.	(3marks)

SECTION B - (40 MARKS)

<u>Answer Question 6 (Compulsory) and Either Question 7 Or 8 in The Spaces Provided After Question 8</u>

6. The relationship between oxygen concentration, sugar consumption and potassium ion uptake in isolated wheat roots was determined. The results obtained were tabulated as shown below. The loss of sugar and potassium uptake or gain are in arbitrary units.

		Perce	ntage ox	ygen in a	erotun st	ream	
	0	5	10	15	20	30	100
Sugar loss	15	20	43	45	45	44	43
Potassium ion gain	5	55	70	75	75	72	70

	ot g me				f s	uga	r le	SSC	and	d p	ota	ass	siuı	m i	on	s ga	ain	ag	gain	st	OX	yge	en	COI	nc	ent	rati	ion	on
		ПП	TII	ПП	711	11111			TITI			П	777	m					TIII			TI	TT		TT				Ш
										-																			
																	1												
							i			1							1					Н	1			Ш			
										1											-								
			Ш		Ш													H											
										i								Ħ				Ш							
			Ш		Ш									Ш			1												
										4							1	Ħ		Ħ		H				Ш			
																	1			i									
					Ш								Ш		Ш					-					H	Ħ			
			Ш																						Ц	Ш			
				-														1	HH	1									
							1			1																			
										H					Н		1	#		1					1				
				Ш	Ш					1				Ш			1												
										i								#											
																										Ш			
			Ш		Ш				H	1						H						H				Ш			
		Ш		Ш				111										4							1				
										i						Ш		Ħ		-	111				H	H			
										1											Ш	Ш							
						Ш																			#		1		
				Щ	Ш		Ш			1				Ш															
									ans																			nrk) arks	
•••				•••	•••																	•••							
									 1d p			 sin	 ım	ion								•••							
									atio		LU DE)1 u		101		>~***										(2	ma	arks	s)
																											· • • •		
											• • •																. 		
				• • •																· • •									

((ii) Bet	ween 5% and 20% oxygen concentration	(2 marks)
• • • • •	• • • • • • • • • • • • • • • • • • • •		
• • • • •	• • • • • • • • • • • • • • • • • • • •		
d) S	Sugges	t two factors necessary for the above process apart from o	
			(2 marks)
e) S	State tv	vo ways by which the process above can be stopped.	(2 marks)
			· ·
	Vame t	wo main areas in a mammalian body where the above pro	cess occurs.
f) N	· varric t	wo main areas in a manimanan body where the above pro	
t) ſ		main areas in a manimanan body where the above pro	(2 marks)
t) r			(2 marks)
			(2 marks)
			(2 marks)
	ain the		(2 marks) Dersal. (20 marks)
Expla	ain the	various ways in which seeds and fruits are adapted to disp scribe how the digestion of a protein is achieved in the foll mentary canal.	(2 marks) Dersal. (20 marks)
Expla (a)	ain the Des alin (i) Sto	various ways in which seeds and fruits are adapted to disp scribe how the digestion of a protein is achieved in the foll mentary canal.	(2 marks) persal. (20 marks) lowing portions of the
Expla (a)	ain the Des alin (i) Sto	various ways in which seeds and fruits are adapted to disp scribe how the digestion of a protein is achieved in the foll mentary canal.	(2 marks) persal. (20 marks) dowing portions of the (4 marks) (4 marks)
Expla (a)	Des alin (i) Stor	various ways in which seeds and fruits are adapted to disposeribe how the digestion of a protein is achieved in the follonentary canal. mach odenum	(2 marks) Dersal. (20 marks) Owing portions of the (4 marks) (4 marks) (4 marks) xylem of the root. (8 marks)
Expla (a)	Des alin (i) Stor (ii) Duo (i)	various ways in which seeds and fruits are adapted to disposeribe how the digestion of a protein is achieved in the follonentary canal. mach odenum Describe the process of absorption at the root hair to the	(2 marks) persal. (20 marks) dowing portions of the (4 marks) (4 marks) (4 marks) xylem of the root. (8 marks) rate of transpiration.
Expla (a)	Des alin (i) Stor (ii) Duo (i)	various ways in which seeds and fruits are adapted to disposeribe how the digestion of a protein is achieved in the follonentary canal. mach odenum Describe the process of absorption at the root hair to the	(2 marks) persal. (20 marks) dowing portions of the (4 marks) (4 marks) (4 marks) xylem of the root. (8 marks) rate of transpiration.
Expla (a)	Des alin (i) Stor (ii) Duo (i)	various ways in which seeds and fruits are adapted to disposeribe how the digestion of a protein is achieved in the follonentary canal. mach odenum Describe the process of absorption at the root hair to the	(2 marks) persal. (20 marks) dowing portions of the (4 marks) (4 marks) (4 marks) xylem of the root. (8 marks) rate of transpiration.
Expla (a)	Des alin (i) Stor (ii) Duo (i)	various ways in which seeds and fruits are adapted to disposeribe how the digestion of a protein is achieved in the follonentary canal. mach odenum Describe the process of absorption at the root hair to the	(2 marks) persal. (20 marks) dowing portions of the (4 marks) (4 marks) (4 marks) xylem of the root. (8 marks) rate of transpiration.
Expla (a)	Des alin (i) Stor (ii) Duo (i)	various ways in which seeds and fruits are adapted to disposeribe how the digestion of a protein is achieved in the follonentary canal. mach odenum Describe the process of absorption at the root hair to the	(2 marks) persal. (20 marks) dowing portions of the (4 marks) (4 marks) (4 marks) xylem of the root. (8 marks) rate of transpiration.
Expla (a)	Des alin (i) Stor (ii) Duo (i)	various ways in which seeds and fruits are adapted to disposeribe how the digestion of a protein is achieved in the follonentary canal. mach odenum Describe the process of absorption at the root hair to the	(2 marks) persal. (20 marks) dowing portions of the (4 marks) (4 marks) (4 marks) xylem of the root. (8 marks) rate of transpiration.

